
CENTER FOR DRUG EVALUATION AND RESEARCH APPROVAL PACKAGE FOR:

APPLICATION NUMBER 21-307

Final Printed Labeling



LOTRIMIN butenafine hydrochloride cream 1%	BUITTATM ANTIFUNGAL	Ne
Active ingradient Butenafine hydrochloride 1%	Purpose	
Uses cures most athlete's foot between the relieves itching, burning, cracking, and	toes, jock itch and ringworm d scaling which accompany these condition	ıs
Warnings For external use only		
Do not use on nails or scalp in or near the mou	th or the eyes • for vaginal yeast infection	s
When using this product do not get into	the eyes. If eye contact occurs, rinse thorou	igh

23564-00

Wt 24g (0.85 oz)

ly with water.

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a dector if too much irritation occurs or gets worse.

Keep out of reach of children. If swallowed, get medicat help or contact a Poison Control Center right away.

Directions

adults and children 12 years and older

use the tip of the cap to break the seal and open the tube

wash the affected skin with soap and water and dry completely before applying

"tor athlets toole between the foes: apply to affected skin between and around the toes twice a day for 1 week (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventitated shoes. Change shoes and socks at least once daily.

Tor jost the ame ringuem apply once a day to affected skin for 2 weeks or as directed by a doctor

wash hands after each use

children under 12 years: ask a doctor


Store at 5° -30°C (41° - 86°F). See crimp for lot number and expiration date

©2001 Distributed by Schering-Plough HealthCare Products, Inc., Memphis, TN 38151 U.S.A. All rights reserved. Made in U.S.A.



LOTRIMINE!	de ann	23565-0
LOTRIMING butenafine hydrochloride cream 1%	U CI " ANTII	UNGAL
tive ingredient tenafine hydrochloride 1%	Perpose	Net Wt 12g (0.42 oz
g res most jock itch lieves itching, burning, cracking, and scaling		condition
l'arnings er external use entr		
on naituse on naits or scalp in or near the mouth or the	avec a fee vaninal wase	1 infactions
When using this product do not get into the eyes		
Hop use ned ask a declar if too much irritation o		
Koop out of reach of children. If swallowed, get med	fical help or contact a Pois	on Control Center right away
Directions		
adults and children 12 years and older	n the tube	
 use the tip of the cap to break the seal and ope wash the affected skin with soap and water an 	d dry completely before :	polvina
apply once a day to affected skin for 2 weeks or	r as directed by a doctor	****
• wash hands affer each use • children under 12 years; ask a doctor		
Store at 5° - 30°C (41° - 86°F). See crimp f	or lot number and eval	ration data

LOTPIMINE.	29562-00	Τ
LOTRIMINE Ultra TM ANTIFUNGAL	Net Wt 15g (0.53 oz)	ı
Active ingratient Propose Butengline hydrochloride 1%		
Uses: a cures most athlete's foot between the toes, jock rich and ringworm a relieves itching, burning, cracking, and scaling which accompany these conditions		1
Warnings For external use only		
Do not use won nails or scalp win or near the mouth or the eyes wfor vaninal yeast i	intections	1
When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly	y with water.	1
Step use and ask a dector if too much irritation occurs or gets worse.		ı
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Cents	fas right away.	1
Directions a dath and children 12 years and older a sole to the cup to break the seal and open the tube a use the tip of the cup to break the seal and open the tube a was the affected skin with soop and water and dry completely before applying the affected is the between the elect apply to affected skin between and around the too (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear welf- Change shows and socks at least once daily. for jeck titch and ringineum apply once a day to affected skin for 2 weeks or as directed a wash hands after each use a children under 12 years; ask a doctor	l-fitting, verrifated shoes.	
Store at 5° - 30°C (41° - 86°F). See crimp for lot number and expiration date of Schering-Plough HealthCare Products, Inc., Memohis, TN 38151 U.S.A. All rights reserve	© 2001 Distributed by red. Made in U.S.A.	



əsodınd

Active ingredient

L MBTOL butenafine hydrochloride cream 1

Drug Facts



CURES MOST JOCK ITCH

- ·Relieves Itching, Burning and Chafing
- •Full Prescription
 Strength

LOTRIMIN

butenafine hydrochloride cream 1% ANTIFUNGAL

LOTRIMIN

butenafine hydrochloride cream 1% ANTIFUNGAL

Net Wt 15g (0.53 oz)

ionore entrettime else su tisiv



esodina

Putensfine hydrochloride cresm 1% butensuren.

TOUSINIA



sinaibarpni avilonil

noilemioini 19110

, 100 , 100


Reep out of reach

Springs Warnings sol ying asu ton ed

Drug Facis Active ingredient

DEFENDENCE PRODUCTION OF THE PROPERTY OF THE P







LOTRIMIN It was but enafine hydrochloride cream 1%.

Drug Facts

LOTRIMI *** 29560-00 butenafine hydrochloride cream 1%** ANTIFUNGAL Net Wr 30g (1.1 oz)	ľ
butenafine hydrochloride cream 1%	=
Active Ingradient Purpose Butenafine hydrochloride 1%	
Uses - cures most athlete's foot between the toes, jock itch and ringworm - relieves itching, burning, cracking, and scaling which accompany these conditions	
Warnings For external use only	
Do not use on nails or scalp in or near the mouth or the eyes of or vaginal yeast infections	·
When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.	· [
Stop use and ask a dector if too much irritation occurs or gets worse.	·
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	·
Directions	·
adults and children 12 years and older	
was the tip of the cap to break the seal and open the tube	
 wash the affected skin with soap and water and dry completely before applying for albiets's foot between the loes; apply to affected skin between and around the toes twice a day for 1 week (morning) 	
and munit, or once a day for 4 weeks, or as directed by a doctor, wear well-litting, ventilated shoes. Change shoes and	
socks at least once daily.	
 for jock lich and ringworm apply once a day to affected skin for 2 weeks or as directed by a doctor wash hands after each use 	
= children under 12 years: ask a doctor	
Store at 5° - 30°C (41° - 86°F). See crimp for lot number and expiration date ©2001 Distributed by Schering-Plough HealthCare Products, Inc., Memphis, TN 38151 U.S.A. All rights reserved. Made in U.S.A.	